



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let m and n be 4 and 2; or 27 and 1; or 9 and 3; etc.; then $2^3 + 1^2 = 3^2$; $3^3 + 13^2 = 14^2$; $3^3 + 3^2 = 6^2$; etc.

V. Solution by H. C. WILKES, Skull Run, West Virginia.

$x^3 = (a+y)(a-y)$. Let $a+y=x^2$ and $a-y=x$, then $x^2+x=2a$, and $x = \frac{1}{2} \pm \sqrt{2a + \frac{1}{4}}$. Let a be any triangular number, and from the above formula, integral values for x , a , and y can be found.

VI. Solution by O. W. ANTHONY, M. Sc., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

Let $x=ky$. Then $x^3+y^2=a^2$ becomes $y^2\{k^3y+1\}=a^2$. This will be a square if $y=k^3+2$. $\therefore y=k^3+2$, and $x=k(k^3+2)$ will be a solution, where k is any integer. If $k=1$, $y=3$, $x=3$ and $x^3+y^2=36$. If $k=2$, $y=10$, $x=20$, and $x^3+y^2=8100$, etc., etc.

VII. Solution by J. H. DRUMMOND, LL. D., Portland, Maine.

(A). If the problem is to be taken literally, $y = \sqrt{a^2 - x^3}$ in which x may any number whose third power $<$ than a^2 . But this does not give exact results.

(B). If it means that $x^3+y^2=\square$, let $x=my$ and we have $m^3y+1=\square=(\text{say}) b^2$ and $y=(b^2-1)/m^3$ and $x=(b^2-1)/m^2$; but then $a=b(b^2-1)/m^3$, in which m and b may be any numbers greater than unity, but the value of a depends on x and y .

(C). By transposing $x^3=a^2-y^2$; take $x=a-y$, then $x^2=a+y$, and $a^2-2ay+y^2=a+y$, and $y=(2a+1 \pm \sqrt{8a+1})/2$. As y must be less than a to make x positive, the sign of the radical term must be negative. It is readily seen that $a=n(n+1)/2$ makes $8a+1$ a square, and by reducing we get $y=n(n-1)/2$ and $x=n$, in which n may be any number.

(D). If the question means to find exact values of x and y for any value of a , I cannot solve it.

46. Proposed by JOSIAH H. DRUMMOND, LL. D., Portland, Maine.

In $x^2 + x\sqrt{xy} = a \dots (1)$ and $y^2 + y\sqrt{xy} = b \dots (2)$ find such values of a and b as will make x and y integral; give a general solution.

I. Solution by the PROPOSER.

Take $y=m^2x$, and by combining the two equations and reducing we have, $\frac{b}{a}(m+1)=m^3(m+1)$ and consequently $m^3=\frac{b}{a}$.

From (1) we have $x=\pm\sqrt{\frac{a}{m+1}}$. Take $a=c^2$ and $m+1=d^2$ and substituting, we have $x=c/d$. To make this value integral, take $c=de$; then $x=e$, and $y=m^2x=e(d^2-1)^2$. But $a=c^2$, and $c=dx=de$. $\therefore a=d^2e^2$; but $b=am^3=d^2e^2(d^2-1)^3$, in which a may be any whole number >1 , and e any whole number.

II. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

In order that $\sqrt[4]{xy}$ be integral and rational, we put $x=rm^2$ and $y=rn^2$, r , m , and n being any integers. Whence we readily find that when $a=r^2m^3(m+n)$ and $b=r^2n^3(m+n)$, x and y are integral.

Now put $r=1$, $m=3$, and $n=2$, and we obtain $x^2+x\sqrt[4]{xy}=135$ and $y^2+y\sqrt[4]{xy}=40$; whence $x=9$ and $y=4$.

Put $r=2$, $m=2$, and $n=1$; then $x^2+x\sqrt[4]{xy}=96$ and $y^2+y\sqrt[4]{xy}=12$; whence $x=8$, and $y=2$.

III. Solution by A. H. BELL, Box 184, Hillsboro, Illinois.

The only condition to fill is to make $xy=\square$. Take $x=4$, $y=1$, and $a=24$, $b=3$, etc., etc.

IV. Solution by H. C. WILKES, Skull Run, West Virginia.

Let $m^2=x$, $n^2=y$. Then $m^3(m+n)=a$; $n^3(m+n)=b$. \therefore To make x and y integral, a and b must have a common factor $(m+n)$. The remaining factors will be m^3 and n^3 . Let $a=448$, $b=189$; then $x=16$, $y=9$. $7(64)m=4$; $7(27)n=3$.

V. Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

Let $P=x^{\frac{1}{2}}$, $Q=y^{\frac{1}{2}}$. Then $P^4+P^3Q=a$ (1). $Q^4+Q^3P=b$ (2).

$$(1) \div (2), P=Q\sqrt[4]{a/b}. \quad \therefore P^2=x=\pm \frac{a^{\frac{1}{2}}}{\sqrt[4]{a^{\frac{1}{2}}+b^{\frac{1}{2}}}}, \quad Q=y=\pm \frac{b^{\frac{1}{2}}}{\sqrt[4]{a^{\frac{1}{2}}+b^{\frac{1}{2}}}.$$

$$\text{Let } a=\{\frac{1}{2}(m^2+n^2)\}^3, \quad b=\{\frac{1}{2}(m^2-n^2)\}^3.$$

$$\therefore x=\pm \frac{(m^2+n^2)^2}{4m}, \quad y=\pm \frac{(m^2-n^2)^2}{4m}.$$

$$\text{Let } m=pn. \quad \therefore x=\pm \frac{n^3(p^2+1)^2}{4p}, \quad y=\pm \frac{n^3(p^2-1)^2}{4p}.$$

$$\text{Let } n=2p. \quad \therefore x=\pm 2p^2(p^2+1)^2, \quad y=\pm 2p^2(p^2-1)^2. \\ \therefore a=\{2p^2(p^2+1)\}^3, \quad b=\{2p^2(p^2-1)\}^3.$$

VI. Solution by O. W. ANTHONY, M. Sc., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland

Let $y=m^2x$. Then $x^2(1+m)=a$, and $x^2m^3(1+m)=b$.

$$\therefore m=\frac{b^{\frac{1}{2}}}{a^{\frac{1}{2}}}. \quad x=\frac{a^{\frac{1}{2}}}{\sqrt[4]{a^{\frac{1}{2}}+b^{\frac{1}{2}}}}; \quad y=\frac{b^{\frac{1}{2}}}{\sqrt[4]{a^{\frac{1}{2}}+b^{\frac{1}{2}}}.$$

$$\text{Let } a=p^3; \quad b=q^3. \quad \text{Then } x=\frac{p^2}{\sqrt[4]{p+q}}; \quad y=\frac{q^2}{\sqrt[4]{p+q}}.$$

Let $p=2rs$; $q=r^2+s^2$. Then $x=\frac{4r^2s^2}{r+s}$; $y=\frac{(r^2+s^2)^2}{r+s}$.

Let $r=k+l$; $s=k-l$. Then $x=\frac{2(k^2-l^2)^2}{k}$; $y=\frac{2(k^2+l^2)^2}{k}$.

Let $l=\alpha k$. Then $x=2k^3(1-\alpha^2)^2$; $y=2k^3(1+\alpha^2)^2$.

Now $a=p^3=8r^3s^3=8(k^2-l^2)^3=8k^6(1-\alpha^2)^3$, and $b=q^3=(r^2+s^2)^3=8(k^2+l^2)^3=8k^6(1+\alpha^2)^3$, where α and k are integers.

PROBLEMS.

53. Proposed by A. H. BELL, Box 184, Hillsboro, Illinois.

Given $x^2-114xy=\mp 3$ to find the least values of x and y in integers.

54. Proposed by JOSIAH H. DRUMMOND, LL. D., Portland, Maine.

In the expression $2x^2-2ax+b^2$, find two series of values for x in integral terms of a and b .

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

35. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

Find the chance that the distance of two points within a square shall not exceed a side of the square. [From *Byerly's Integral Calculus*.]

I. Solution by ALWYN C. SMITH, The University of Colorado, Boulder, Colorado.

a is one side of the square; P and Q the two points; (x, y) the point P with O for origin; and r and ϕ the polar coordinates of Q , with P as origin. Then the favorable cases are

$$4 \int_0^{\frac{1}{2}\pi} \int_0^a \int_0^{a-r\sin\phi} \int_0^{a-r\cos\phi} dx dy r dr d\phi = a^4(\pi - \frac{1}{6}).$$

All the cases $= a^2 \cdot a^2 = a^4$. Therefore, $p = \pi - \frac{1}{6}$.

